



10071


A2




(1-1)




(1-2)



(1-3)



(1-4)




(1-5)

**IN THE CLAIMS:**

Please cancel claims 1-44

Please enter new claims 45-54:


45. A method for treating an inflammatory respiratory disease comprising administering to a subject a therapeutically effective amount of a vitamin D<sub>3</sub> derivative described by the following general formula (3) or pharmaceutically permissible solvate thereof,



wherein, R<sub>01</sub> and R<sub>02</sub> are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyltrimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group; the configuration of the carbon atom at the 20-position is (R)-


configuration;  $R_7$  is a methyl group of a methylene group; when  $R_7$  is a methylene group, the bond between  $R_7$  and the carbon atom at the 26-position is double bond.

46. A method for treating an inflammatory respiratory disease comprising administering to a subject a therapeutically effective amount of a vitamin D<sub>3</sub> derivative expressed by the following general formula (1) or pharmaceutically permissible solvates thereof,



{wherein,  $R_{01}$  and  $R_{02}$  are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyldimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group,

Z is one out of the following formulae (1-1), (1-2), (1-3), (1-4) or (1-5),



[in the above formulae (1-1) to (1-5),

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

m is an integer of 0 to 2;

n is an integer of 0 to 2;

X' is an oxygen atom or NH;

R<sub>11</sub> and R<sub>12</sub> are identical to or different from each other, and express a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group;

K, L and M are each a hydrogen atom; M is a hydrogen atom, and K and L together express a single bond and express a double bond in cooperation with the single bond already shown in the formula; or K is a hydrogen atom, and L and M together express a single bond and express a double bond in cooperation with the single bond already shown in the formula;

R<sub>21</sub>, R<sub>22</sub> and R<sub>23</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxy group, a carboxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>1</sub>-C<sub>4</sub> alkyloxycarbonyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or R<sub>21</sub> and R<sub>22</sub> together may express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

Q expresses >C(-F)-R<sub>31</sub> or >N-R<sub>31</sub>, and herein R<sub>31</sub> is a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxy group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>32</sub>, R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

A and B are identical to or different from each other, and they express a hydrogen atom or a hydroxyl group, or together express a single bond and form a double bond in cooperation with the single bond already shown in the formula;

X and Y together express a carbonyl group in cooperation with the carbon atom to which they are bonded, one of them is a hydrogen atom and the other is a hydroxyl group, or one of them is a hydrogen atom and the other is a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

R<sub>41</sub> and R<sub>42</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or they express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>43</sub> and R<sub>44</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>45</sub> and R<sub>46</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

D and E express each a hydrogen atom, D is a hydroxy group and E expresses a hydrogen atom, D and E together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, or E and R<sub>41</sub> together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, wherein D expresses a hydrogen atom or a hydroxy group; and R<sub>42</sub> expresses a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;


R<sub>51</sub> expresses -CONR<sub>511</sub>R<sub>512</sub>, -COR<sub>513</sub> or -C(OH)R<sub>514</sub>R<sub>515</sub>, wherein R<sub>511</sub> and R<sub>512</sub> are identical to or different from each other, and they are a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group, or both the members together express a nitrogen-containing C<sub>3</sub>-C<sub>8</sub> alkyl ring or a morpholino group in cooperation with the nitrogen atom to which they are bonded; and R<sub>513</sub>, R<sub>514</sub> and R<sub>515</sub> are identical to or different from each other, and they express a C<sub>1</sub>-C<sub>4</sub> alkyl group;

R<sub>52</sub> expresses a methyl group, an ethyl group, a trifluoromethyl group or a pentafluoroethyl group,]

with the proviso that the following compounds (a), (b) and (c) are excluded,


(a) a compound in which the groups of one combination out of R<sub>21</sub> and R<sub>22</sub>, R<sub>32</sub> and R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub>, R<sub>41</sub> and R<sub>42</sub>, R<sub>43</sub> and R<sub>44</sub>, and R<sub>45</sub> and R<sub>46</sub> are both hydroxy groups, both alkyloxy groups, or a hydroxy group and an alkyloxy group,

(b) a compound expressed by the above formula (1) in which Z is the following formula (1-6),



(wherein, p and q are each 0 or the integer 1;  $R_6$  is a hydrogen atom or a  $C_1$ - $C_4$  alkyl group), and

(c) a compound of the following formula (2),



(wherein,  $R_{01}$  and  $R_{02}$  are defined in the same manner as in the above formula (1); the configuration of the carbon atom at the 20-position is (R)-configuration;  $R_7$  is a methyl group or a methylene group; when  $R_7$  is a methylene group, the bond between  $R_7$  and the carbon atom at the 26-position is double bond)).


47. A treating agent for an inflammatory respiratory disease described in Claim 45 or 46, wherein the inflammatory respiratory disease is one or not less than two kinds of inflammatory respiratory diseases selected from a group consisting of acute upper airway infection, chronic sinusitis, allergic rhinitis, chronic lower airway infection, pulmonary emphysema, pneumonia, bronchial asthma, tuberculosis sequela, acute respiratory distress syndrome, cystic fibrosis and pulmonary fibrosis.

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

48. A treating agent for an inflammatory respiratory disease described in Claim 47, wherein the acute upper airway infection is one or not less than two kinds of diseases selected from a group consisting of common cold, acute pharyngitis, acute rhinitis, acute sinusitis, acute tonsillitis, acute pharyngitis, acute epiglottitis and acute bronchitis.

49. A treating agent for an inflammatory respiratory disease described in Claim 47, wherein the chronic lower airway infection is one or not less than two kinds of diseases selected from a group consisting of chronic bronchitis, diffuse panbronchiolitis and bronchiectasis.


50. A method for treating one or not less than two kinds of inflammatory respiratory diseases selected from a group consisting of chronic bronchitis, diffuse panbronchiolitis, bronchiectasis, bronchial asthma, pulmonary emphysema, tuberculosis sequela and cystic fibrosis comprising administering to a subject a therapeutically effective amount of a vitamin D<sub>3</sub> derivative expressed by the following formula (1) or pharmaceutically permissible solvate thereof:



Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

{wherein,  $R_{01}$  and  $R_{02}$  are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyldimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group;

Z is one out of the following formulae (1-1), (1-2), (1-3), (1-4) or (1-5),



[in the above formulae (1-1) to (1-5),

$m$  is an integer of 0 to 2;

$n$  is an integer of 0 to 2;

$X'$  is an oxygen atom or NH;

$R_{11}$  and  $R_{12}$  are identical to or different from each other, and express a hydrogen atom or a  $C_1$ - $C_4$  alkyl group;

$K$ ,  $L$  and  $M$  are each a hydrogen atom;  $M$  is a hydrogen atom, and  $K$  and  $L$  together express a single bond and express a double bond in cooperation with the single bond already shown in the formula; or  $K$  is a hydrogen atom, and  $L$  and  $M$  together express a single bond and express a double bond in cooperation with the single bond already shown in the formula;

$R_{21}$ ,  $R_{22}$  and  $R_{23}$  are identical to or different from each other, and they are a hydrogen atom, a hydroxy group, a carboxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_1$ - $C_4$  alkyloxycarbonyl group, a  $C_2$ - $C_5$  acyloxy group, a  $C_1$ - $C_4$  alkyloxy group or a  $C_1$ - $C_4$  alkyl

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or R<sub>21</sub> and R<sub>22</sub> together may express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

Q expresses >C(-F)-R<sub>31</sub> or >N-R<sub>31</sub>, and herein R<sub>31</sub> is a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxy group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>32</sub>, R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

A and B are identical to or different from each other, and they express a hydrogen atom or a hydroxyl group, or together express a single bond and form a double bond in cooperation with the single bond already shown in the formula;

X and Y together express a carbonyl group in cooperation with the carbon atom to which they are bonded, one of them is a hydrogen atom and the other is a hydroxyl group, or one of them is a hydrogen atom and the other is a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

R<sub>41</sub> and R<sub>42</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or they express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

*Handwritten: CB*

R<sub>43</sub> and R<sub>44</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

*Handwritten: CB*

R<sub>45</sub> and R<sub>46</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

*Handwritten: CB*

D and E express each a hydrogen atom, D is a hydroxy group and E expresses a hydrogen atom, D and E together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, or E and R<sub>41</sub> together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, wherein D expresses a hydrogen atom or a hydroxy group; and R<sub>42</sub> expresses a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>51</sub> expresses -CONR<sub>511</sub>R<sub>512</sub>, -COR<sub>513</sub> or -C(OH)R<sub>514</sub>R<sub>515</sub>, wherein R<sub>511</sub> and R<sub>512</sub> are identical to or different from each other, and they are a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group, or both the members together express a nitrogen-containing C<sub>3</sub>-C<sub>8</sub> alkyl ring or a morpholino group

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167


in cooperation with the nitrogen atom to which they are bonded; and  $R_{513}$ ,  $R_{514}$  and  $R_{515}$  are identical to or different from each other, and they express a  $C_1$ - $C_4$  alkyl group;

$R_{52}$  expresses a methyl group, an ethyl group, a trifluoromethyl group or a pentafluoroethyl group,]

with the proviso that the following compounds (a), (b) and (c) are excluded,

(a) a compound in which the groups of one combination out of  $R_{21}$  and  $R_{22}$ ,  $R_{32}$  and  $R_{33}$ ,  $R_{34}$  and  $R_{35}$ ,  $R_{41}$  and  $R_{42}$ ,  $R_{43}$  and  $R_{44}$ , and  $R_{45}$  and  $R_{46}$  are both hydroxy groups, both alkyloxy groups, or a hydroxy-group and an alkyloxy group,


(b) a compound expressed by the above formula (1) in which Z is the following formula (1-6),



(1-6)

(wherein,  $p$  and  $q$  are each 0 or the integer 1;  $R_6$  is a hydrogen atom or a  $C_1$ - $C_4$  alkyl group), and

(c) a compound of the following formula (2),




(2)

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

(wherein,  $R_{01}$  and  $R_{02}$  are defined in the same manner as in the above formula (1); the configuration of the carbon atom at the 20-position is (R)-configuration;  $R_7$  is a methyl group or a methylene group; when  $R_7$  is a methylene group, the bond between  $R_7$  and the carbon atom at the 26-position is double bond)).


51. A method for treating a disease selected from a group consisting of malignant tumors, rheumatoid arthritis, osteoporosis, diabetes mellitus, hypertension, alopecia, acne, psoriasis and dermatitis comprising administering to a subject a therapeutically effective amount of a vitamin D<sub>3</sub> derivative expressed by the following formula (1) or pharmaceutically permissible solvate thereof:



{wherein,  $R_{01}$  and  $R_{02}$  are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyl dimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group;

Z is one out of the following formulae (1-1), (1-2), (1-3), (1-4) or (1-5),

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167



[in the above formulae (1-1) to (1-5),

m is an integer of 0 to 2;

n is an integer of 0 to 2;

X' is an oxygen atom or NH;

R<sub>11</sub> and R<sub>12</sub> are identical to or different from each other, and express a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group;

K, L and M are each a hydrogen atom; M is a hydrogen atom, and K and L together express a single bond and express a double bond in cooperation with the single bond already shown in the formula; or K is a hydrogen atom, and L and M together express a single bond and express a double bond in cooperation with the single bond already shown in the formula;

R<sub>21</sub>, R<sub>22</sub> and R<sub>23</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxy group, a carboxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>1</sub>-C<sub>4</sub> alkyloxycarbonyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or R<sub>21</sub> and R<sub>22</sub> together may express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

Q expresses  $>C(-F)-R_{31}$  or  $>N-R_{31}$ , and herein  $R_{31}$  is a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_2-C_5$  acyloxy group, a  $C_1-C_4$  alkyloxy group or a  $C_1-C_4$  alkyl group which may be substituted with a hydroxy group, a  $C_2-C_5$  acyloxy group or a  $C_1-C_4$  alkyloxy group;

$R_{32}$ ,  $R_{33}$ ,  $R_{34}$  and  $R_{35}$  are identical to or different from each other, and they are a hydrogen atom, a hydroxyl group, a  $C_1-C_4$  alkyl group or a  $C_2-C_5$  acyloxy group;

A and B are identical to or different from each other, and they express a hydrogen atom or a hydroxyl group, or together express a single bond and form a double bond in cooperation with the single bond already shown in the formula;

X and Y together express a carbonyl group in cooperation with the carbon atom to which they are bonded, one of them is a hydrogen atom and the other is a hydroxyl group, or one of them is a hydrogen atom and the other is a  $C_2-C_5$  acyloxy group;

$R_{41}$  and  $R_{42}$  are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_2-C_5$  acyloxy group, a  $C_1-C_4$  alkyloxy group or a  $C_1-C_4$  alkyl group which may be substituted with a hydroxyl group, a  $C_2-C_5$  acyloxy group or a  $C_1-C_4$  alkyloxy group, or both the members together express a  $C_1-C_5$  alkylidene group, or they express a  $C_3-C_6$  cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

$R_{43}$  and  $R_{44}$  are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_2-C_5$  acyloxy group, a  $C_1-C_4$  alkyloxy group or a  $C_1-C_4$  alkyl group which may be substituted with a hydroxyl group, a  $C_2-C_5$  acyloxy group or a  $C_1-C_4$  alkyloxy group, or both the members together express a

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

C<sub>1</sub>-C<sub>5</sub> alkylidene group, or express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>45</sub> and R<sub>46</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

D and E express each a hydrogen atom, D is a hydroxy group and E expresses a hydrogen atom, D and E together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, or E and R<sub>41</sub> together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, wherein D expresses a hydrogen atom or a hydroxy group; and R<sub>42</sub> expresses a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>51</sub> expresses -CONR<sub>511</sub>R<sub>512</sub>, -COR<sub>513</sub> or -C(OH)R<sub>514</sub>R<sub>515</sub>, wherein R<sub>511</sub> and R<sub>512</sub> are identical to or different from each other, and they are a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group, or both the members together express a nitrogen-containing C<sub>3</sub>-C<sub>8</sub> alkyl ring or a morpholino group in cooperation with the nitrogen atom to which they are bonded; and R<sub>513</sub>, R<sub>514</sub> and R<sub>515</sub> are identical to or different from each other, and they express a C<sub>1</sub>-C<sub>4</sub> alkyl group;


R<sub>52</sub> expresses a methyl group, an ethyl group, a trifluoromethyl group or a pentafluoroethyl group;

with the proviso that the following compounds (a), (b) and (c) are excluded,

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

(a) a compound in which the groups of one combination out of  $R_{21}$  and  $R_{22}$ ,  $R_{32}$  and  $R_{33}$ ,  $R_{34}$  and  $R_{35}$ ,  $R_{41}$  and  $R_{42}$ ,  $R_{43}$  and  $R_{44}$ , and  $R_{45}$  and  $R_{46}$  are both hydroxy groups, both alkyloxy groups, or a hydroxy group and an alkyloxy group,


(b) a compound expressed by the above formula (1) in which Z is the following formula (1-6),



(1-6)

(wherein, p and q are each 0 or the integer 1; R<sub>6</sub> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), and


(c) a compound of the following formula (2),



(2)


(wherein, R<sub>01</sub> and R<sub>02</sub> are defined in the same manner as in the above formula (1); the configuration of the carbon atom at the 20-position is (R)-configuration; R<sub>7</sub> is a methyl group or a methylene group; when R<sub>7</sub> is a methylene group, the bond between R<sub>7</sub> and the carbon atom at the 26-position is double bond)).

52. A method for treating hypercalcemia attributable to vitamin D excess comprising administering to a subject a therapeutically effective amount of a vitamin D<sub>3</sub> derivative of the following Formula (1) or pharmaceutically permissible solvate thereof,



{wherein, R<sub>01</sub> and R<sub>02</sub> are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyltrimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group;

Z is one out of the following formulae (1-2), (1-3), (1-4) or (1-5),



[in the above formulae (1-2) to (1-5),

m is an integer of 0 to 2;

n is an integer of 0 to 2;

X' is an oxygen atom or NH;

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

*Handwritten: B1*

$R_{21}$ ,  $R_{22}$  and  $R_{23}$  are identical to or different from each other, and they are a hydrogen atom, a hydroxy group, a carboxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_1$ - $C_4$  alkyloxycarbonyl group, a  $C_2$ - $C_5$  acyloxy group, a  $C_1$ - $C_4$  alkyloxy group or a  $C_1$ - $C_4$  alkyl group which may be substituted with a hydroxyl group, a  $C_2$ - $C_5$  acyloxy group or a  $C_1$ - $C_4$  alkyloxy group, or  $R_{21}$  and  $R_{22}$  together may express a  $C_3$ - $C_6$  cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

*Handwritten: A2*

Q expresses  $>C(-F)-R_{31}$  or  $>N-R_{31}$ , and herein  $R_{31}$  is a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_2$ - $C_5$  acyloxy group, a  $C_1$ - $C_4$  alkyloxy group or a  $C_1$ - $C_4$  alkyl group which may be substituted with a hydroxy group, a  $C_2$ - $C_5$  acyloxy group or a  $C_1$ - $C_4$  alkyloxy group;

$R_{32}$ ,  $R_{33}$ ,  $R_{34}$  and  $R_{35}$  are identical to or different from each other, and they are a hydrogen atom, a hydroxyl group, a  $C_1$ - $C_4$  alkyl group or a  $C_2$ - $C_5$  acyloxy group;

A and B are identical to or different from each other, and they express a hydrogen atom or a hydroxyl group, or together express a single bond and form a double bond in cooperation with the single bond already shown in the formula;

X and Y together express a carbonyl group in cooperation with the carbon atom to which they are bonded, one of them is a hydrogen atom and the other is a hydroxyl group, or one of them is a hydrogen atom and the other is a  $C_2$ - $C_5$  acyloxy group;

$R_{41}$  and  $R_{42}$  are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_2$ - $C_5$  acyloxy group, a  $C_1$ - $C_4$  alkyloxy group or a  $C_1$ - $C_4$  alkyl group which may be substituted with a hydroxyl group, a  $C_2$ - $C_5$  acyloxy group or a  $C_1$ - $C_4$  alkyloxy group, or both the members together express a

C<sub>1</sub>-C<sub>5</sub> alkylidene group, or they express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>43</sub> and R<sub>44</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>45</sub> and R<sub>46</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

D and E express each a hydrogen atom, D is a hydroxy group and E expresses a hydrogen atom, D and E together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, or E and R<sub>41</sub> together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, wherein D expresses a hydrogen atom or a hydroxy group; and R<sub>42</sub> expresses a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>51</sub> expresses -CONR<sub>511</sub>R<sub>512</sub>, -COR<sub>513</sub> or -C(OH)R<sub>514</sub>R<sub>515</sub>, wherein R<sub>511</sub> and R<sub>512</sub> are identical to or different from each other, and they are a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group, or

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167


both the members together express a nitrogen-containing C<sub>3</sub>-C<sub>8</sub> alkyl ring or a morpholino group in cooperation with the nitrogen atom to which they are bonded; and R<sub>513</sub>, R<sub>514</sub> and R<sub>515</sub> are identical to or different from each other, and they express a C<sub>1</sub>-C<sub>4</sub> alkyl group;

R<sub>52</sub> expresses a methyl group, an ethyl group, a trifluoromethyl group or a pentafluoroethyl group,]

with the proviso that the following compounds (a), (b) and (c) are excluded,

(a) a compound in which the groups of one combination out of R<sub>21</sub> and R<sub>22</sub>, R<sub>32</sub> and R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub>, R<sub>41</sub> and R<sub>42</sub>, R<sub>43</sub> and R<sub>44</sub>, and R<sub>45</sub> and R<sub>46</sub> are both hydroxy groups, both alkyloxy groups, or a hydroxy group and an alkyloxy group,


(b) a compound expressed by the above formula (1) in which Z is the following formula (1-6),



(1-6)


(wherein, p and q are each 0 or the integer 1; R<sub>6</sub> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), and

(c) a compound of the following formula (2),




(wherein,  $R_{01}$  and  $R_{02}$  are defined in the same manner as in the above formula (1); the configuration of the carbon atom at the 20-position is (R)-configuration;  $R_7$  is a methyl group or a methylene group; when  $R_7$  is a methylene group, the bond between  $R_7$  and the carbon atom at the 26-position is double bond)).

53. A method for treating hypoparathyroidism comprising administering to a subject a therapeutically effective amount of a vitamin  $D_3$  derivative of the following Formula (1) or pharmaceutically permissible solvate thereof,



{wherein,  $R_{01}$  and  $R_{02}$  are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyl dimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group;

Z is one out of the following formulae (1-2), (1-3), (1-4) or (1-5),



[in the above formulae (1-2) to (1-5),

m is an integer of 0 to 2;

n is an integer of 0 to 2;

X' is an oxygen atom or NH;

R<sub>21</sub>, R<sub>22</sub> and R<sub>23</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxy group, a carboxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>1</sub>-C<sub>4</sub> alkyloxycarbonyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or R<sub>21</sub> and R<sub>22</sub> together may express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

Q expresses >C(-F)-R<sub>31</sub> or >N-R<sub>31</sub>, and herein R<sub>31</sub> is a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxy group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>32</sub>, R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

A and B are identical to or different from each other, and they express a hydrogen atom or a hydroxyl group, or together express a single bond and form a double bond in cooperation with the single bond already shown in the formula;

X and Y together express a carbonyl group in cooperation with the carbon atom to which they are bonded, one of them is a hydrogen atom and the other is a hydroxyl group, or one of them is a hydrogen atom and the other is a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

R<sub>41</sub> and R<sub>42</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or they express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>43</sub> and R<sub>44</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>45</sub> and R<sub>46</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

B1  
A3  
D and E express each a hydrogen atom, D is a hydroxy group and E expresses a hydrogen atom, D and E together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, or E and R<sub>41</sub> together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, wherein D expresses a hydrogen atom or a hydroxy group; and R<sub>42</sub> expresses a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;


R<sub>51</sub> expresses -CONR<sub>511</sub>R<sub>512</sub>, -COR<sub>513</sub> or -C(OH) R<sub>514</sub>R<sub>515</sub>, wherein R<sub>511</sub> and R<sub>512</sub> are identical to or different from each other, and they are a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group, or both the members together express a nitrogen-containing C<sub>3</sub>-C<sub>8</sub> alkyl ring or a morpholino group in cooperation with the nitrogen atom to which they are bonded; and R<sub>513</sub>, R<sub>514</sub> and R<sub>515</sub> are identical to or different from each other, and they express a C<sub>1</sub>-C<sub>4</sub> alkyl group;

R<sub>52</sub> expresses a methyl group, an ethyl group, a trifluoromethyl group or a pentafluoroethyl group,]

with the proviso that the following compounds (a), (b) and (c) are excluded,

(a) a compound in which the groups of one combination out of R<sub>21</sub> and R<sub>22</sub>, R<sub>32</sub> and R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub>, R<sub>41</sub> and R<sub>42</sub>, R<sub>43</sub> and R<sub>44</sub>, and R<sub>45</sub> and R<sub>46</sub> are both hydroxy groups, both alkyloxy groups, or a hydroxy group and an alkyloxy group,


(b) a compound expressed by the above formula (1) in which Z is the following formula (1-6),



(1-6)

(wherein, p and q are each 0 or the integer 1; R<sub>6</sub> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), and


(c) a compound of the following formula (2),



(2)





(wherein, R<sub>01</sub> and R<sub>02</sub> are defined in the same manner as in the above formula (1); the configuration of the carbon atom at the 20-position is (R)-configuration; R<sub>7</sub> is a methyl group or a methylene group; when R<sub>7</sub> is a methylene group, the bond between R<sub>7</sub> and the carbon atom at the 26-position is double bond)).

54. A method for treating metabolic disorder of cartilage comprising administering to a subject a therapeutically effective amount of a vitamin D<sub>3</sub> derivative of the following Formula (1) or pharmaceutically permissible solvate thereof,



{wherein,  $R_{01}$  and  $R_{02}$  are each independently a hydrogen atom, a trimethylsilyl group, a triethylsilyl group, a t-butyldimethylsilyl group, an acetyl group, a methoxymethyl group or a tetrahydro-4H-pyran-2-yl group;

Z is one out of the following formulae (1-2), (1-3), (1-4) or (1-5),



[in the above formulae (1-2) to (1-5),

m is an integer of 0 to 2;

n is an integer of 0 to 2;

$X'$  is an oxygen atom or NH;

$R_{21}$ ,  $R_{22}$  and  $R_{23}$  are identical to or different from each other, and they are a hydrogen atom, a hydroxy group, a carboxyl group, a trifluoromethyl group, a pentafluoroethyl group, a  $C_1$ - $C_4$  alkyloxycarbonyl group, a  $C_2$ - $C_5$  acyloxy group, a  $C_1$ - $C_4$  alkyloxy group or a  $C_1$ - $C_4$  alkyl group which may be substituted with a hydroxyl group, a  $C_2$ - $C_5$  acyloxy group or a  $C_1$ - $C_4$

alkyloxy group, or R<sub>21</sub> and R<sub>22</sub> together may express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

Q expresses >C(-F)-R<sub>31</sub> or >N-R<sub>31</sub>, and herein R<sub>31</sub> is a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxy group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>32</sub>, R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub> are identical to or different from each other, and they are a hydrogen atom, a hydroxyl group, a C<sub>1</sub>-C<sub>4</sub> alkyl group or a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

A and B are identical to or different from each other, and they express a hydrogen atom or a hydroxyl group, or together express a single bond and form a double bond in cooperation with the single bond already shown in the formula;

X and Y together express a carbonyl group in cooperation with the carbon atom to which they are bonded, one of them is a hydrogen atom and the other is a hydroxyl group, or one of them is a hydrogen atom and the other is a C<sub>2</sub>-C<sub>5</sub> acyloxy group;

R<sub>41</sub> and R<sub>42</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or they express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>43</sub> and R<sub>44</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy

Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group, or both the members together express a C<sub>1</sub>-C<sub>5</sub> alkylidene group, or express a C<sub>3</sub>-C<sub>6</sub> cyclic alkyl group in cooperation with the carbon atom to which they are bonded;

R<sub>45</sub> and R<sub>46</sub> are identical to or different from each other, and they express a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

D and E express each a hydrogen atom, D is a hydroxy group and E expresses a hydrogen atom, D and E together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, or E and R<sub>41</sub> together express a single bond and express a double bond in cooperation with the single bond already shown in the formula, wherein D expresses a hydrogen atom or a hydroxy group; and R<sub>42</sub> expresses a hydrogen atom, a hydroxyl group, a trifluoromethyl group, a pentafluoroethyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group, a C<sub>1</sub>-C<sub>4</sub> alkyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyl group which may be substituted with a hydroxyl group, a C<sub>2</sub>-C<sub>5</sub> acyloxy group or a C<sub>1</sub>-C<sub>4</sub> alkyloxy group;

R<sub>51</sub> expresses -CONR<sub>511</sub>R<sub>512</sub>, -COR<sub>513</sub> or -C(OH)R<sub>514</sub>R<sub>515</sub>, wherein R<sub>511</sub> and R<sub>512</sub> are identical to or different from each other, and they are a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group, or both the members together express a nitrogen-containing C<sub>3</sub>-C<sub>8</sub> alkyl ring or a morpholino group in cooperation with the nitrogen atom to which they are bonded; and R<sub>513</sub>, R<sub>514</sub> and R<sub>515</sub> are identical to or different from each other, and they express a C<sub>1</sub>-C<sub>4</sub> alkyl group;


Preliminary Amendment  
Divisional of U. S. Patent No. 09/830,167

R<sub>52</sub> expresses a methyl group, an ethyl group, a trifluoromethyl group or a pentafluoroethyl group,]

with the proviso that the following compounds (a), (b) and (c) are excluded,

(a) a compound in which the groups of one combination out of R<sub>21</sub> and R<sub>22</sub>, R<sub>32</sub> and R<sub>33</sub>, R<sub>34</sub> and R<sub>35</sub>, R<sub>41</sub> and R<sub>42</sub>, R<sub>43</sub> and R<sub>44</sub>, and R<sub>45</sub> and R<sub>46</sub> are both hydroxy groups, both alkyloxy groups, or a hydroxy group and an alkyloxy group,


(b) a compound expressed by the above formula (1) in which Z is the following formula (1-6),



(1-6)

(wherein, p and q are each 0 or the integer 1; R<sub>6</sub> is a hydrogen atom or a C<sub>1</sub>-C<sub>4</sub> alkyl group), and

(c) a compound of the following formula (2),



(2)

(wherein, R<sub>01</sub> and R<sub>02</sub> are defined in the same manner as in the above formula (1); the configuration of the carbon atom at the 20-position is (R)-configuration; R<sub>7</sub> is a methyl group or